- and management. *Dermatol Ther* 2006;19:202-209.
4. Lucky AW, Biro FM, Huster GA, Leach AD, Morrison JA, Ratterman J. Acne vulgaris in premenarchal girls. An early sign of puberty associated with rising levels of dehydroepiandrosterone. *Arch Dermatol* 1994;130:308-314.
 5. Lucky AW. A review of infantile and pediatric acne. *Dermatology* 1998;196:95-97.
 6. Plewig G, Kligman AM. Acne and rosacea. 2nd ed. Berlin: Springer-Verlag, 1993:674-675.
 7. Sardana K, Sharma RC, Sarkar R. Seasonal variation in acne vulgaris—myth or reality. *J Dermatol* 2002;29:484-488.
 8. Hancox JG, Sheridan SC, Feldman SR, Fleischer AB Jr. Seasonal variation of dermatologic disease in the USA: a study of office visits from 1990 to 1998. *Int J Dermatol* 2004;43:6-11.
 9. Karciauskienė J, Valiukeviciene S, Gollnick H, Stang A. The prevalence and risk factors of adolescent acne among schoolchildren in Lithuania: a cross-sectional study. *J Eur Acad Dermatol Venereol* 2014;28:733-740.
 10. Thiboutot D, Del Rosso JQ. Acne vulgaris and the epidermal barrier: is acne vulgaris associated with inherent epidermal abnormalities that cause impairment of barrier functions? Do any topical acne therapies alter the structural and/or functional integrity of the epidermal barrier? *J Clin Aesthet Dermatol* 2013;6:18-24.

<https://doi.org/10.5021/ad.2019.31.2.251>

Squamous Cell Carcinoma on the Fingers of Orthopedic Surgeon Induced by Occupational Radiation Exposure

Hee Seong Yoon, Ji Hye Heo, Si Hyub Lee, Jeonghyun Shin, Gwang Seong Choi, Ji Won Byun

Department of Dermatology, Inha University School of Medicine, Incheon, Korea

Dear Editor:

Recently, fluoroscopy is useful for diagnosis and treatment and its usage is increasing in the treatment of pain, orthopedic surgery and interventional radiology. The occupational radiation exposure of physicians is also increasing and the hand is the most exposed area. Radiodermatitis of physicians caused by occupational radiation exposure has been reported¹, but there is few report of the development of squamous cell carcinoma (SCC). We describe an orthopedic surgeon with multiple SCCs on the fingers occupationally exposed to radiation.

A 49-year-old man presented with painful crusted necrotic plaques and scale widely distributed in the both fingers 2

years ago (Fig. 1A~E). He smoked 5~6 packs of tobacco every day. He was an orthopedic surgeon and repeatedly exposed to radiation because of frequent use of fluoroscopy. For 11 years, he performed an average of 700 cases of percutaneous vertebral augmentation and 70,000 cases of nerve block per year without radiation shielding gloves. The skin biopsy showed atypical squamous cells invading into dermis forming sheet and individual keratinization (Fig. 2). Magnetic resonance imaging of the right hand suspected bone invasion in the right index finger (Fig. 1H). Chest computed tomography revealed slightly enlarged lymph nodes in both axilla; however, they appeared benign. We diagnosed him with SCC and pre-

Received January 5, 2018, Revised April 2, 2018, Accepted for publication April 15, 2018

Corresponding author: Ji Won Byun, Department of Dermatology, Inha University School of Medicine, 27 Inhang-ro, Jung-gu, Incheon 22332, Korea. Tel: 82-32-890-2238, Fax: 82-32-890-2236, E-mail: piggy1981@hanmail.net
ORCID: <https://orcid.org/0000-0003-0317-6725>

This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/4.0>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

Copyright © The Korean Dermatological Association and The Korean Society for Investigative Dermatology

Fig. 1. Painful crusted necrotic plaques and scale widely distributed in the right thumb, index, and 3rd finger, and left thumb and index finger. Each figure is as follows: (A) Both hands (B) right thumb (C) right index and 3rd finger (D) left thumb (E) left index finger. And photographs after 17 months of follow-up. Each figure is as follows: (F) Dorsal side of the left hand. There were three SCC lesions on the left index finger (arrows). (G) Ventral side of the right hand. There was a SCC lesion on the right third finger (arrow). (H) Right hand MRI image. On the T2 weighted image, SCC invasion is suspected in the medial cortex of the mid phalanx and the marrow of the distal phalanx of the right index finger. SCC: squamous cell carcinoma, MRI: magnetic resonance imaging.

scribed multidisciplinary cancer care. For the right thumb and index finger, the SCC lesions were widely excised. The distal phalanx of the right index finger, which had a high possibility of bone involvement, was partially severed leaving a 6-mm of distal phalanx, and the defect was repaired with a free flap. Mohs micrographic surgery (MMS) and a full thickness skin graft (FTSG) was performed on the left thumb. All fingers suspected of skin cancer in situ due to chronic radiation dermatitis were treated with ingenol mebutate. During 17 months of follow-up, he often wore protective gloves and continued to work. However, new SCC lesions of the right middle, left index finger appeared (Fig. 1F, G). MMS and FTSG was performed for those lesions. He is under observation without recurrence until 25 months later.

The effects of radiation exposure on the skin was described by the International Commission on Radiological

Protection in 2010². The annual equivalent dose is 500 mSv to the skin². Direct beam contact with fluoroscopy causes radiation exposure of 40 mSv per minute, and no deep eye radiation exposure was observed at a distance of 91.4 cm from the fluoroscope³. The skin effects of radiation are divided into acute and chronic changes, depending on the time period from the exposure⁴. Chronic changes include epidermal thinning, fibrosis, edema, dermal thickening, dyspigmentation, telangiectasia, and dermal necrosis⁴. The risk of cutaneous cancer increases after radiation exposure⁴.

Surgical methods are mainly used for the treatment of SCC. Among the surgical methods, MMS has a 96% cure rate and is the gold standard⁵. If there is bone involvement, amputation should be performed⁵. Since cancer has been reported up to 50 years after exposure, continuous follow-up observation is needed¹. There is concern in this case

Fig. 2. (A, B) Histopathologic findings of the nail bed of the right thumb (H&E: A, $\times 40$; B, $\times 200$). The skin biopsy showed atypical squamous cells invading into dermis forming sheet and individual keratinization.

that SCC will recur, so we will continue follow-up. This case demonstrates that physicians should be aware of occupational radiation exposure.

ACKNOWLEDGMENT

We thank the patient for granting permission to publish this information. This work was supported by the Inha University Research Grant and grants (56990) from the National Research Foundation through the Korean government (NRF-2016R1C1B1014905).

CONFLICTS OF INTEREST

The authors have nothing to disclose.

ORCID

Hee Seong Yoon, <https://orcid.org/0000-0001-8997-9697>
 Ji Hye Heo, <https://orcid.org/0000-0002-0928-122X>
 Si Hyub Lee, <https://orcid.org/0000-0002-8246-8664>
 Jeonghyun Shin, <https://orcid.org/0000-0002-4995-9533>

Gwang Seong Choi, <https://orcid.org/0000-0002-5766-0179>
 Ji Won Byun, <https://orcid.org/0000-0003-0317-6725>

REFERENCES

1. Shim DM, Kim YM, Oh SK, Lim CM, Kwon BT. Radiation induced hand necrosis of an orthopaedic surgeon who had treated a patient with fluoroscopy-guided spine injection. *J Korean Orthop Assoc* 2014;49:250-254.
2. ICRP. Radiological protection in fluoroscopically guided procedures outside the imaging department. ICRP Publication 117. *Ann ICRP* 2010;40.
3. Mehlman CT, DiPasquale TG. Radiation exposure to the orthopaedic surgical team during fluoroscopy: "how far away is far enough?". *J Orthop Trauma* 1997;11:392-398.
4. Hegedus F, Mathew LM, Schwartz RA. Radiation dermatitis: an overview. *Int J Dermatol* 2017;56:909-914.
5. Dika E, Fanti PA, Patrizi A, Misciali C, Vaccari S, Piraccini BM. Mohs surgery for squamous cell carcinoma of the nail unit: 10 years of experience. *Dermatol Surg* 2015;41:1015-1019.